


Biomass Coverage in Norway, Iceland, Faroe Islands Region

Biomass Coverage Over Time in Norway, Iceland, Faroe Islands Region

Stocks ordered from lowest to highest average biomass




Taxonomy Group Gadids Pleuronectids Sebastids Forage Fish




Herring ICES 1–2–4a–5–14a (Norwegian Spring Spawners) – Norway, Iceland, Faroe Islands Region Stock represents 24% of average absolute surplus production

Taxonomy Group Forage Fish




Capelin Barents Sea – Norway, Iceland, Faroe Islands Region

Taxonomy Group Forage Fish



Atlantic cod North–East Arctic – Norway, Iceland, Faroe Islands Region Stock represents 17% of average absolute surplus production


Taxonomy Group Gadids




Atlantic cod Iceland Grounds – Norway, Iceland, Faroe Islands Region

Taxonomy Group Gadids

Herring ICES 1–2–4a–5–14a (Norwegian Spring Spawners) in Norway, Iceland, Faroe Islands Region




Capelin Barents Sea in Norway, Iceland, Faroe Islands Region




First and last years with recorded surplus production are labeled

Atlantic cod North–East Arctic in Norway, Iceland, Faroe Islands Region



First and last years with recorded surplus production are labeled

Atlantic cod Iceland Grounds in Norway, Iceland, Faroe Islands Region



First and last years with recorded surplus production are labeled