




Biomass Coverage in Sebastids Taxonomy Group




Biomass Coverage Over Time in Sebastids Taxonomy Group

Stocks with highest average biomass



Biomass Coverage Over Time in Sebastids Taxonomy Group


Stocks with lowest average biomass



Region ■ Canada West Coast ■ Japan ■ US Alaska ■ US West Coast

Deepwater redfish North-East Arctic – Sebastids Taxonomy Group


Stock represents 13% of average absolute surplus production



Region ■ Norway, Iceland, Faroe Islands


Pacific ocean perch Eastern Bering Sea and Aleutian Islands – Sebastids Taxonomy Group

Stock represents 3% of average absolute surplus production




Golden redfish ICES 5–6–12–14 – Sebastids Taxonomy Group

Stock represents 12% of average absolute surplus production




Pacific ocean perch Gulf of Alaska – Sebastids Taxonomy Group

Stock represents 6% of average absolute surplus production




Deepwater redfish North–East Arctic in Sebastids Taxonomy Group




First and last years with recorded surplus production are labeled

Pacific ocean perch Eastern Bering Sea and Aleutian Islands in Sebastids Taxonomy Group




First and last years with recorded surplus production are labeled

Golden redfish ICES 5-6-12-14 in Sebastids Taxonomy Group



First and last years with recorded surplus production are labeled

Pacific ocean perch Gulf of Alaska in Sebastids Taxonomy Group



First and last years with recorded surplus production are labeled